



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Book

Search PubMed for #3 AND #4

☒ Limits Preview/Index History Clipboard Details

About Entrez

Field: Title

Text Version

Show: 20 Sort Text

Items 1-4 of 4

One page

Entrez PubMed
Overview
Help | FAQ
Tutorial
New/Noteworthy
E-Utilities

PubMed Services
Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources
Order Documents
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

Privacy Policy

- ☐ 1: [Deng WG, Saunders MA, Gilroy DW, He XZ, Yeh H, Zhu Y, Shtivelband MI, Ruan KH, Wu KK.](#) Related Articles, Li
Purification and characterization of a cyclooxygenase-2 and angiogenesis suppressing factor produced by human fibroblasts.
FASEB J. 2002 Aug;16(10):1286-8. Epub 2002 Jun 07.
PMID: 12060668 [PubMed - indexed for MEDLINE]
- ☐ 2: [Danz H, Stoyanova S, Wippich P, Brattstrom A, Hamburger M.](#) Related Articles, Li
Identification and isolation of the cyclooxygenase-2 inhibitory principle in Isatis tinctor
Planta Med. 2001 Jul;67(5):411-6.
PMID: 11488453 [PubMed - indexed for MEDLINE]
- ☐ 3: [Abad T, McNaughton-Smith G, Fletcher WQ, Echeverri F, Diaz-Penate R, Tabraue C, Ruiz de Galarreta CM, Lopez-Blanco F, Luis JG.](#) Related Articles, Li
Isolation of (S)-(+)-naproxene from Musa acuminata. Inhibitory effect of naproxene and its 7-methoxy isomer on constitutive COX-1 and inducible COX-2.
Planta Med. 2000 Jun;66(5):471-3.
PMID: 10909271 [PubMed - indexed for MEDLINE]
- ☐ 4: [Percival MD, Ouellet M, Vincent CJ, Yergey JA, Kennedy BP, O'Neill GP.](#) Related Articles, Li
Purification and characterization of recombinant human cyclooxygenase-2.
Arch Biochem Biophys. 1994 Nov 15;315(1):111-8.
PMID: 7979387 [PubMed - indexed for MEDLINE]

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Freedom of Information Act](#) | [Disclaimer](#)

Apr 13 2004 06:



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Book

Search PubMed for [] Go Clear

☒ Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Show: 20 Sort Send to Text

Text Version

☐ 1: Proc Natl Acad Sci U S A. 1992 Aug 15;89(16):7384-8.

Related Articles, L

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

FREE full text article at
www.pnas.org

FREE full text article
in PubMed Central

Human cyclooxygenase-2 cDNA.

Hla T, Neilson K.

Department of Molecular Biology, Holland Laboratory, American Red Cross, Rockville MD 20855.

Cyclooxygenase (Cox), also known as prostaglandin (PG) H synthase (EC 1.14.99.1), catalyzes the rate-limiting step in the formation of inflammatory PGs. A major regulator step in PG biosynthesis is at the level of Cox: growth factors, cytokines, and tumor promoters induce Cox activity. We have cloned the second form of the Cox gene (Cox-2) from human umbilical vein endothelial cells (HUVEC). The cDNA encodes a polypeptide of 604 amino acids that is 61% identical to the previously isolated human Cox-1 polypeptide. In vitro translation of the human (h)Cox-2 transcript in rabbit reticulocyte lysates resulted in the synthesis of a 70-kDa protein that is immunoprecipitated by antiserum to ovine Cox. Expression of the hCox-2 open reading frame in Cos-7 monkey kidney cells results in the elaboration of cyclooxygenase activity. hCox-2 cDNA hybridizes to a 4.5-kilobase mRNA species in HUVEC, whereas the hCox-1 cDNA hybridizes to 3- and 5.3-kilobase species. Both Cox-1 and Cox-2 mRNAs are expressed in HUVEC, vascular smooth muscle cells, monocytes, and fibroblasts. Cox-2 mRNA was preferentially induced by phorbol 12-myristate 13-acetate and lipopolysaccharide in human endothelial cells and monocytes. Together, these data demonstrate that the Cox enzyme is encoded by at least two genes that are expressed and differentially regulated in a variety of cell types. High-level induction of the hCox-2 transcript in mesenchymal-derived inflammatory cells suggests a role in inflammatory conditions.

PMID: 1380156 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to Text

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Freedom of Information Act | Disclaimer

Apr 13 2004 06.

L5 ANSWER 213 OF 213 DRUGU COPYRIGHT 2004 THOMSON DERWENT on STN
ACCESSION NUMBER: 1994-46604 DRUGU B P E
TITLE: **Characterization of the mechanism of inhibition of
human cyclooxygenase-2 by
anti-inflammatory drugs.**
AUTHOR: Ouellet M; Percival M D
CORPORATE SOURCE: Merck-Frosst
LOCATION: Kirkland, Quebec, Canada
SOURCE: Can.J.Physiol.Pharmacol. (72, Suppl. 1, 453, 19 1 Ref.
CODEN: CJPPA3 ISSN: 0008-4212
AVAIL. OF DOC.: Merck Frosst Centre for Therapeutic Research, Kirkland, QC
H9R 4P8, Canada.
LANGUAGE: English
DOCUMENT TYPE: Journal
FIELD AVAIL.: AB; LA; CT
FILE SEGMENT: Literature

AB The kinetic mechanism of inhibition of the **purified** form of
inducible cyclooxygenase (hCox-2) by several classical NSAIDs and a
selective Cox-2 inhibitor was determined. Flurbiprofen, meclofenamate
and indometacin were all time-dependent inhibitors of hCox-2. None of
the 3 inhibitors had a high degree of selectivity, when compared with
hCox-1. NS-398 also had time-dependent inhibition of hCox-2, but was a
time-independent inhibitor of hCox-1. The difference in the mechanism of
inhibition was reflected in the high degree of selectivity observed for
hCox-2 over hCox-1. Results demonstrate that the mechanism of inhibition
of hCox-2 by classical NSAIDs is similar to that identified for ovine
Cox-1. In addition, it shows the nature of time-dependency of inhibition
of hCox-1 and hCox-2 greatly determines the degree of selectivity for 1
isozyme over the other. (conference abstract).